

Review Article

The Role of Zinc in Alzheimer's Disease

Nicole T. Watt, Isobel J. Whitehouse, and Nigel M. Hooper

Institute of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, Clarendon Way, Leeds LS2 9JT, UK

Correspondence should be addressed to Nigel M. Hooper, n.m.hooper@leeds.ac.uk

Received 14 September 2010; Accepted 9 November 2010

Academic Editor: Anthony R. White

Copyright © 2011 Nicole T. Watt et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Zinc, the most abundant trace metal in the brain, has numerous functions, both in health and in disease. Zinc is released into the synaptic cleft of glutamatergic neurons alongside glutamate from where it interacts and modulates NMDA and AMPA receptors. In addition, zinc has multifactorial functions in Alzheimer's disease (AD). Zinc is critical in the enzymatic nonamyloidogenic processing of the amyloid precursor protein (APP) and in the enzymatic degradation of the amyloid- β ($A\beta$) peptide. Zinc binds to $A\beta$ promoting its aggregation into neurotoxic species, and disruption of zinc homeostasis in the brain results in synaptic and memory deficits. Thus, zinc dyshomeostasis may have a critical role to play in the pathogenesis of AD, and the chelation of zinc is a potential therapeutic approach.

1. Introduction

Alzheimer's disease (AD) is the most prevalent form of dementia, which affects more than 37 million people worldwide, with an estimated cost of \$422 billion in 2009 [1, 2]. Moreover, the incidence of the illness and the prospect of an aging population will result in rising social and economic demands. AD is characterised by the presence of extracellular amyloid plaques and intracellular neurofibrillary tangles within the afflicted brain, which cause neuronal loss in the neocortex, hippocampus, and basal forebrain, leading to progressive cognitive and behavioural decline [3]. Zinc, in addition to copper and iron, has been shown to be involved in AD. Here, we review the current literature relating to neuronal zinc metabolism and the way in which zinc can modulate normal brain activity. We discuss also the contribution of zinc to the formation, aggregation, and degradation of the amyloid- β ($A\beta$) peptide and the contribution of zinc to the pathogenesis of AD.

2. Physiological Role of Zinc in the Brain

As the most abundant trace metal in the brain, zinc is found tightly associated with numerous proteins conferring either structural or catalytic properties upon them [4]. However, a significant amount of loosely bound, chelatable zinc can

be found sequestered in presynaptic vesicles forming a sub-population of "zinc enriched" (ZEN) neurones [5, 6], which co-release zinc with the neurotransmitter glutamate upon excitation. The majority of these "gluzincergic" neurones [7] have their cell bodies located in either the cerebral cortex or the limbic structures of the forebrain [8], and so an extensive network uniting limbic and cerebrocortical functions is created [9]. This connection between zinc and glutamatergic neurotransmission allows the ion to modulate the overall excitability of the brain and also influence synaptic plasticity [10].

The identification of synaptic zinc was first made by McLardy over fifty years ago who identified that a band of zinc dithizonate staining correlated with hippocampal mossy fibre axons [7]. Since then, many more cerebrocortical pathways have been identified which contain zinc-rich synaptic vesicles; indeed nearly 50% of the glutamatergic synapses are actually "gluzincergic" in some parts of the cerebral cortex. Significantly, only small amounts of chelatable zinc can be determined in glutamatergic pathways which originate outside the cerebral cortex or limbic systems.

Despite this extensive network of zinc-containing neurons, little is known about how zinc homeostasis is maintained within the neuron. There are two families of zinc transporters: the ZnT family, which act to decrease intracellular zinc concentrations by exporting zinc from the cytoplasm to the lumen of organelles or the extracellular

space, and the ZIP family, which import the metal from the extracellular space or organellar lumen into the cytoplasm [11]. Whilst many of the transporters have particular distribution patterns, only ZnT3 expression is restricted to the brain and the testis [12]. It is located in the vesicular membrane [13] and is necessary to transport zinc from the cytoplasm into the synaptic vesicle of the neuron. The vesicular concentration of zinc correlates with the amount of ZnT3 present [14]. Targeted disruption of ZnT3 in a mouse model resulted in a complete lack of chelatable zinc [15].

A number of approaches have been taken to confirm that zinc is coreleased with glutamate from the presynaptic bouton during neuronal excitation: imaging of zinc in boutons before and after stimulation [16], analytical detection of zinc released into perfusates [17], and direct imaging of released zinc using fluorescent extracellular probes [18, 19]. This latter approach has provided the most definitive results. An early study employed a reporter construct whose fluorescence properties changed upon zinc binding. Stimulation of organotypic cultures from rat hippocampus produced a cloud of green fluorescence as the released zinc bound an apometalloenzyme confirming the release of zinc from the culture [17]. A later study by Quinta-Ferreira and colleagues [20] demonstrated a release of zinc with each pulse of an action potential. Whilst there is now no doubt that zinc is released during synaptic activity, there is little consensus on the amount or duration of its existence in the synaptic cleft [21].

Following an intense burst of neuronal activity, the release of glutamate and postsynaptic membrane depolarisation open a variety of zinc-permeable ion channels which contribute to removing the ion from the extracellular fluid. These include N-methyl-D-aspartate (NMDA) channels and calcium permeable α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA)/kainate channels. The consequences of zinc acting on these receptors are diverse and demonstrate the significance zinc has in modulating fast excitatory glutamatergic transmission. Zinc can act to either enhance or depress synaptic activity with varying degrees of potency [21].

The most studied interaction is zinc with NMDA receptors (NMDAR). Initially, zinc was thought to selectively inhibit NMDAR-mediated neuronal activity by inducing a voltage-independent noncompetitive inhibition that decreased the probability of the channel being open [22–24]. A voltage-dependent inhibition of NMDAR could be observed at higher concentrations of zinc ($>20 \mu\text{M}$) and was believed to be due to binding of the cation within the channel pore [25]. With the cloning of NMDAR subunits, it was confirmed that zinc could cause both voltage-independent and voltage-dependent inhibition [26]. The exceptional sensitivity of the GluN2A subunit towards zinc suggests that even contaminating ions found in routine laboratory solutions are sufficient to cause inhibition [27]. Significantly, despite being responsible for inhibitory effects at NMDAR, it has also been shown that NMDAR activation may provide a route of influx for zinc contributing to toxic effects of exposure [28] (Figure 1).




FIGURE 1: Multiple mechanisms for zinc uptake following synaptic release. Zinc and glutamate are released from a “gluzincergic” synapse during neurotransmission. The actions of glutamate, alongside membrane depolarisation, open a number of zinc-permeable channels on the postsynaptic membrane which clear the ion from the extracellular fluid. (1) AMPA receptors; (2) NMDA receptors; (3) voltage-gated calcium channels; (4) TRPM7. (blue pentagons) zinc ion and (green ellipses) glutamate molecule. For simplicity, zinc-permeable channels are only shown on the post-synaptic membrane.

Whilst it is thought that zinc exposure generally attenuates NMDAR-mediated neurotoxicity, zinc has been shown to potentiate AMPAR-mediated toxicity at suggested physiological concentrations ($50 \mu\text{M}$). Originally, it was proposed that the toxic effect was due to zinc influx via voltage-gated calcium channels [29], with Lin et al. [30] showing that desensitisation of AMPAR would explain such an enhancement. Subsequently, it has been demonstrated that zinc can carry currents directly via AMPAR, mostly via the calcium-permeable subtype [31] (Figure 1). At high supraphysiological doses (1 mM), zinc has been shown to inhibit AMPAR [21, 32]. A few studies have also looked directly at zinc-mediated inhibition of voltage-gated calcium channels, as their proximity to vesicular release sites on the presynaptic membrane suggests they could interact [33, 34]. Most recently, neurotoxicity has been attributed to transient receptor potential metastasin 7 (TRPM7) channel activation resulting in increased intracellular zinc [35] (Figure 1).

The significance of synaptically released zinc centres on the amount that is released into the synaptic cleft upon excitation. Some authors argue that the amount of zinc ($10\text{--}100 \mu\text{M}$) released following an action potential arriving at the glutamatergic bouton is high enough to bring about the voltage-dependent inhibition of NMDAR and that there would be no spillover onto neighbouring cells [36, 37]. Other authors disagree, suggesting that the zinc concentration would be sufficient to affect nearby cells [38]. It has also been demonstrated that the zinc level (low nM) is only sufficient to block the voltage-independent component of NMDAR activity [39]. Alternatively, it could be that there is little or no diffusible zinc released, supporting the notion that zinc behaves in a “tonic” mode. Kay and Tóth [40] proposed that zinc is exocytosed from the presynaptic membrane and that instead of diffusing into the extracellular space it remains tightly bound to an as-yet unidentified presynaptic component. This would create a “vener” [41] of zinc ions

which would build up with synaptic activity or erode with quiescence to modulate plasticity.

Thus, the implication is that there are three different groups of zinc signals. First, “synaptic zinc” acts as a conventional neurotransmitter, is contained within presynaptic vesicles, and is released upon excitation and binds to a variety of receptors on the postsynaptic membrane. The downstream consequence of receptor binding is one of tonic modulation of glutamatergic excitatory synapses. The second type is similar to calcium signalling and occurs in conjunction with synaptic zinc signalling. A transmembrane flux of “synaptic zinc” from the extracellular space passes through post-synaptic zinc-permeable ion channels. The third is “intracellular zinc signalling” whereby existing intracellular stores are released [7]. This class is difficult to define and as yet has not been identified in neurons but has been demonstrated in mast cells [42].

Therefore, zinc can be classified as an endogenous modulator of synaptic transmission. It is found in synaptic vesicles, released upon excitation, and has multiple synaptic targets. The significant inhibitory effect of zinc on NMDAR, alongside the crucial function of NMDAR in both neurophysiology and pathophysiology, advocates a vital role for zinc in both healthy and diseased brains [21].

3. Role of Zinc in APP Processing

The most prominent lesions in the brain of AD sufferers are the amyloid or “senile” plaques, which predominantly consist of $A\beta$ peptides derived from the proteolytic processing of the amyloid precursor protein (APP). APP is an ubiquitously expressed glycosylated transmembrane protein with a large N-terminal extracellular domain, a single hydrophobic transmembrane domain, and a small C-terminal cytoplasmic domain. A specific and saturable binding site for zinc ($K_D = 750$ nM) has been reported in the cysteine-rich region on the ectodomain of APP [44, 45]. It is hypothesised that zinc could have a role in sustaining the adhesiveness of APP during cell-cell and cell-matrix interactions [46, 47]. APP can be processed by one of two pathways: the amyloidogenic pathway, leading to the production of $A\beta$, and the non-amyloidogenic pathway (Figure 2(a)), reviewed in [48].

In the amyloidogenic pathway, APP is sequentially cleaved by the aspartyl protease, β -site APP-cleaving enzyme 1 (BACE1) forming the secreted $APP\beta$ (sAPP β) fragment and a membrane bound C-terminal fragment of 99 amino acids (C99). The C99 fragment is then further processed by the γ -secretase complex into APP intracellular domain (AICD) and $A\beta$ peptides, predominantly 40 and 42 amino acids in length. It is these aggregation-prone $A\beta$ peptides which form oligomeric and fibrillar structures which deposit in the brain and over time cause AD. The γ -secretase complex comprises four components: presenilin (PS) 1 or 2, nicastrin (Nct), presenilin enhancer 2 (PEN-2), and anterior pharynx defective-1 (Aph-1) [49]. sAPP β is involved in the pruning of synapses during the development of central and peripheral neurons [50] and AICD is known to be a transcription

factor for several genes, including the upregulation of the $A\beta$ -degrading enzyme, neprilysin (NEP) [51].

The predominant APP-processing pathway in healthy brain is the nonamyloidogenic pathway where APP is cleaved by the α -secretase within the $A\beta$ region forming the secreted $APP\alpha$ (sAPP α) fragment and the membrane-bound C-terminal fragment of 83 amino acids (C83) (Figure 2(a)). C83 is subsequently cleaved by the γ -secretase complex generating AICD and p3 (Figure 2(a)). α -secretase activity is attributed to the disintegrin and metalloprotease (ADAM) family of zinc metalloproteases (reviewed in [48, 52]). The ADAMs, along with the matrix metalloproteases (MMPs), are members of the metzincin clan of metalloproteases as they have a long zinc-binding consensus sequence *HEBXHXBGBXH* (H, histidine, zinc ligand; E, catalytic glutamate; G, glycine; B, bulky apolar residue; X, any amino acid) which contains three zinc ligands [53]. Structurally, the catalytic domain is globular and divided into two subdomains, with the active site cleft running between the two [54]. The defining feature of the metzincins is the conserved methionine residue which creates a 1,4- β -turn (Met-turn), creating the catalytic cleft [53]. The catalytic zinc atom sits at the bottom of the groove between the subdomains, with the subsites in the groove determining specificity for particular amino acid sequences in the substrate (reviewed in [52]). A number of enzymes in this family, namely, ADAM9, 10, 17 (also known as TNF- α converting enzyme, TACE) and 19, are known to exert α -secretase activity, but it is unclear which enzyme or enzymes are responsible for the α -cleavage of APP *in vivo* [55–58]. ADAM10 appears to be the predominant enzyme as overexpression of functional ADAM10 in AD transgenic mice led to an increase in sAPP α and reduced $A\beta$ production, plaque deposition, and cognitive deficits [59]. Although fibroblasts from ADAM10^{-/-} mice showed no deficiency in α -secretase activity, probably due to compensation by ADAM17 [60], a recent detailed study has provided strong evidence that ADAM10 is the physiologically relevant constitutive α -secretase in primary neurons [61].

4. Role of Zinc in $A\beta$ Degradation

In a healthy brain, the relatively small amount of $A\beta$ -constitutively being produced is rendered safe by $A\beta$ degrading enzymes. The steady state levels of $A\beta$ synthesis and clearance in cerebrospinal fluid (CSF) are 7.6% and 8.3% per hour, respectively [62]. A large number of candidate $A\beta$ -degrading enzymes have been identified, with the majority being zinc metalloproteases. These include NEP, insulin-degrading enzyme (IDE), endothelin-converting enzyme (ECE) 1 and 2, MMP2, 3, and 9, and PreP (reviewed in [43]). NEP, ECE1 and 2, and MMPs have the conserved zinc-binding motif *HEXXH* (H, histidine, zinc ligand; E, catalytic glutamate; X, any amino acid), and IDE and its homolog PreP are both inverzincins, as they contain the inverted zinc-binding motif *HXXEH*.

NEP appears to be the dominant $A\beta$ protease [63–66], and is capable of degrading monomeric and oligomeric $A\beta$ [63, 67] (Figure 2(b)). NEP knockout mice have significantly

Ser 8 [93], Glu11 [90], and Tyr10 [95] (Figure 3) however, Tyr10 has been ruled out [89] and Arg5 has been deemed highly unlikely [91]. The carboxyl side chain of Glu11 is a zinc ligand [88, 90] however, Asp1 is considered the most attractive zinc ligand, either through its N-terminal amino group and/or its side-chain carboxylate group [87, 88, 91, 93]. Raman spectroscopy has shown zinc binding to the N τ site of histidine side chains in senile plaques taken from AD brains [98], but it is unclear if zinc only binds to free A β peptides that subsequently aggregate or whether zinc binds to A β in preformed plaques.

The reported apparent binding constants (a Kd) of zinc to A β peptides vary from 1 to 300 μ M (reviewed in [91]). The published a Kd values vary greatly due to the *in vitro* conditions (e.g., buffer composition, pH), A β fragment, and experimental method. The high a Kd (20–300 μ M) values come from tyrosine fluorescence experiments however, even these are contentious and hard to reproduce [99–101], and any change in tyrosine fluorescence could be due to A β aggregation rather than zinc binding [101]. Discounting the tyrosine fluorescence measurements, the most likely apparent a Kd value for zinc binding to A β peptides is a range of 1–20 μ M [91]. The binding affinity of zinc for preformed A β fibrils is approximately the same as of the peptides with a a Kd of 1–20 μ M [89, 101].

6. Role of Zinc in Alzheimer's Disease

Numerous studies have looked to address whether zinc levels change with AD progression. It has been shown that there is a significant increase in serum [102] and hippocampal [5] zinc in AD patients compared to age-matched controls. Jiménez-Jiménez et al. [103] demonstrated a significant decrease in CSF zinc but could find no difference in serum zinc levels between AD and age-matched controls. A decrease in serum zinc has also been reported, though it is possible that some of the AD patients included in one study were malnourished [104, 105]. Overall, there is currently no consensus on what happens to zinc concentrations in AD subjects though much of the discrepancy could be put down to differences in patient allocation, sample type, postmortem interval, or type of analysis used.

Alternatively, a redistribution of zinc could be sufficient to promote disease progression. Lovell and coworkers [106, 107] have mapped the expression levels of a number of the ZnT zinc transporters in AD. ZnT-1, 4, and 6 were all found to show increased expression in early stages of disease, though ZnT-1 expression was decreased during mild cognitive impairment [106, 107]. Although it is unknown whether increased expression necessarily correlates with increased activity, these changes in transporter level could result in modified subcellular zinc concentrations. An increase in ZnT6 would lead to an increase in zinc in the TGN [108] which could reduce α -secretase activity [83].

It is well established that amyloid plaques contain increased concentrations of copper, iron, and zinc [98, 109]. Whilst copper and iron appear to be primarily responsible for the toxicity of A β via oxidative-stress-type mechanisms [109, 110], zinc has a crucial role in A β aggregation which is

the most well-established contribution that zinc may have in AD pathogenesis. Whilst the concentration of zinc required for fibrillisation to occur is contentious, with concentrations differing by 100-fold being suggested, zinc is an unequivocal partner in the process [83, 111, 112]. In 2006, Dong and co-workers were able to show that zinc could control the rate of self-assembly of the A β peptides and go on to regulate the amyloid morphology via specific coordination sites [113]. Furthermore, it has been demonstrated that zinc can spontaneously coordinate both intra- and inter-molecular bridging between two peptides to promote A β aggregation [114] and that synaptic zinc promotes A β oligomer formation and their accumulation at excitatory synapses [115].

Studies with synthetic A β showed that chelation chemistry could help solubilise amyloid plaques, with depletion of zinc having a more marked effect on extracting A β than depletion of copper [116]. Oral treatment with 5-chloro-7-iodo-8-hydroxyquinoline (Clioquinol CQ) in Tg2576 mice resulted in a 49% reduction in cortical amyloid deposition [117]. Although CQ has a fairly low affinity (nM) for both copper and zinc, it was still able to release the ions from the A β binding site [118]. A pilot phase II trial in humans showed a decrease in cognitive decline and a reduction in plasma A β_{1-42} in moderately severe AD compared with placebo control [119]. It has been suggested that although CQ may chelate copper and zinc from metallated A β and promote disaggregation, it may not completely halt the aggregation process [120]. A second generation chelator (PBT-2), with improved blood brain barrier penetration, has just completed a phase II clinical trial in early AD with promising results showing good tolerance, a reduction in CSF A β and neuropsychological testing [121].

Recently, it has been shown that zinc can also accelerate the aggregation of a Tau peptide under reducing conditions [122]. Zinc inhibited the formation of intramolecular disulphide bonds but promoted intermolecular bonds between key cysteine residues. Furthermore, zinc exposure has been shown to increase the phosphorylation of PI3K and MAPK-dependent pathways which are key players in Tau modifications [123].

The essential requirement for ZnT3 in loading zinc into synaptic vesicles would suggest that this transporter could have a major impact on zinc signalling in the neuron, even regulating cognitive function. Whilst a lack of zinc signalling in brain slices from ZnT3 $^{-/-}$ mice confirmed the vesicular origin of the released zinc, the mice failed to express a cognitive phenotype. Initial studies detailed a 20% reduction in total zinc level and a loss of histochemically reactive zinc in the synaptic vesicle; however, there was no impairment of spatial learning, memory, or sensorimotor function [124]. The implication being that the vesicular zinc is not required for cognitive function or that compensatory mechanisms made up for the deficits. However, a follow-up study demonstrated marked differences in learning and memory when an older (6 month) cohort of mice was used [125], suggesting that the lack of effect in the previous study was due to the young age (6–10 weeks) of the mice and highlights the importance of aging (the most significant risk

factor) when modelling AD pathology. The results obtained from the older ZnT3^{-/-} cohort established a requirement for zinc in memory function and the maintenance of synaptic health upon aging. Adlard and colleagues proposed that β -amyloid pathology could cause cognitive impairment by trapping zinc within plaques rather than via a directly toxic mechanism [125]. The zinc immobilisation by amyloid would then have similar consequences as a loss of ZnT3 activity with a loss of zinc-dependent synaptic modulation promoting cognitive decline

An alternative approach to minimising the consequences of released zinc could be to promote mechanisms which enhance reuptake. Recently, it has been shown that the cellular form of the prion protein (PrP^C) is an evolutionary descendent of the ZIP family of divalent metal transporters. In particular, ZIPs 5, 6, and 10 were found to have a "prion-like" domain with significant structural similarity. As both PrP^C and the ZIPs bind divalent metal ions via histidine-rich motifs contained within N-terminal repeating sequences, this could suggest that PrP^C is involved in zinc sensing, scavenging, or transport [126]. In agreement with that possibility, we have shown that PrP^C promotes zinc uptake (Watt et al., unpublished). Ensuring efficient clearance of extracellular zinc from the synaptic cleft via PrP^C would exploit an existing physiological process. Furthermore, enhancing zinc uptake would help prevent its ability to contribute to the synaptic targeting of A β oligomers, thus preserving synaptic function [115] and maintaining the proposed ferroxidase activity of APP [127]. As PrP^C levels decrease with age and in sporadic AD [128], it is possible that zinc is cleared less efficiently from the synaptic cleft enhancing aggregation of A β and inhibiting APP ferroxidase activity to promote a pro-oxidative environment. This would suggest that preserving PrP^C function during AD could provide multifactorial benefits, an inhibition of BACE1 which would reduce A β formation [129] and ensure efficient clearance of zinc from the synaptic cleft to prevent aggregation of A β peptides, as well as provide protection against oxidative stress [130, 131].

7. Conclusions

It is clear that zinc not only plays critical roles in the structural and functional integrity of many proteins, but that it also modulates the activity of glutamatergic synapses and indeed may act as a neurotransmitter in its own right. Several of the enzymes involved in processing APP and A β are zinc metalloproteases, with an essential requirement for zinc in their catalytic activity. Zinc binds to A β , promoting its aggregation and thereby modulating its neurotoxicity. Although zinc dyshomeostasis may contribute to the development of AD, further work is required to clarify the molecular and cellular mechanisms affected by zinc under both normal and disease situations.

Acknowledgments

The authors gratefully acknowledge the financial support of the Medical Research Council. N. T. Watt and I. J. Whitehouse contributed equally to this work.

References

- [1] C. Mount and C. Downton, "Alzheimer disease: progress or profit?" *Nature Medicine*, vol. 12, no. 7, pp. 780–784, 2006.
- [2] A. Wimo, B. Winblad, and L. Jönsson, "The worldwide societal costs of dementia: estimates for 2009," *Alzheimer's and Dementia*, vol. 6, no. 2, pp. 98–103, 2010.
- [3] D. J. Selkoe, "Alzheimer's disease: genes, proteins, and therapy," *Physiological Reviews*, vol. 81, no. 2, pp. 741–766, 2001.
- [4] N. M. Hooper, "The biological roles of zinc and families of zinc metalloproteases," in *Zinc Metalloproteases in Health and Disease*, N. M. Hooper, Ed., pp. 1–21, Talor & Francis, London, UK, 1996.
- [5] G. Danscher, K. B. Jensen, C. J. Frederickson et al., "Increased amount of zinc in the hippocampus and amygdala of Alzheimer's diseased brains: a proton-induced X-ray emission spectroscopic analysis of cryostat sections from autopsy material," *Journal of Neuroscience Methods*, vol. 76, no. 1, pp. 53–59, 1997.
- [6] C. J. Frederickson and G. Danscher, "Zinc-containing neurons in hippocampus and related CNS structures," *Progress in Brain Research*, vol. 83, pp. 71–84, 1990.
- [7] C. J. Frederickson and A. I. Bush, "Synaptically released zinc: physiological functions and pathological effects," *BioMetals*, vol. 14, no. 3–4, pp. 353–366, 2001.
- [8] L. Slomianka, G. Danscher, and C. J. Frederickson, "Labeling of the neurons of origin of zinc-containing pathways by intraperitoneal injections of sodium selenite," *Neuroscience*, vol. 38, no. 3, pp. 843–854, 1990.
- [9] J. Y. Koh, "Endogenous zinc in neurological diseases," *Journal of Clinical Neurology*, vol. 1, pp. 121–133, 2005.
- [10] Y. Tian, Z. Yang, and T. Zhang, "Zinc ion as modulator effects on excitability and synaptic transmission in hippocampal CA1 neurons in Wistar rats," *Neuroscience Research*, vol. 68, no. 3, pp. 167–175, 2010.
- [11] L. A. Lichten and R. J. Cousins, "Mammalian zinc transporters: nutritional and physiologic regulation," *Annual Review of Nutrition*, vol. 29, pp. 153–176, 2009.
- [12] R. J. McMahon and R. J. Cousins, "Mammalian zinc transporters," *Journal of Nutrition*, vol. 128, no. 4, pp. 667–670, 1998.
- [13] R. D. Palmiter, T. B. Cole, C. J. Quaipe, and S. D. Findley, "ZnT-3, a putative transporter of zinc into synaptic vesicles," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 93, no. 25, pp. 14934–14939, 1996.
- [14] D. H. Linkous, J. M. Flinn, J. Y. Koh et al., "Evidence that the ZNT3 protein controls the total amount of elemental zinc in synaptic vesicles," *Journal of Histochemistry and Cytochemistry*, vol. 56, no. 1, pp. 3–6, 2008.
- [15] T. B. Cole, H. J. Wenzel, K. E. Kafer, P. A. Schwartzkroin, and R. D. Palmiter, "Elimination of zinc from synaptic vesicles in the intact mouse brain by disruption of the ZnT3 gene," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 4, pp. 1716–1721, 1999.
- [16] Y. Li, C. J. Hough, S. W. Suh, J. M. Sarvey, and C. J. Frederickson, "Rapid translocation of Zn²⁺ from presynaptic terminals into postsynaptic hippocampal neurons after physiological stimulation," *Journal of Neurophysiology*, vol. 86, no. 5, pp. 2597–2604, 2001.
- [17] R. B. Thompson, W. O. Whetsell Jr., B. P. Maliwal, C. A. Fierke, and C. J. Frederickson, "Fluorescence microscopy of stimulated Zn(II) release from organotypic cultures of mammalian hippocampus using a carbonic anhydrase-based

- biosensor system," *Journal of Neuroscience Methods*, vol. 96, no. 1, pp. 35–45, 2000.
- [18] C. J. Frederickson, J. Y. Koh, and A. I. Bush, "The neurobiology of zinc in health and disease," *Nature Reviews Neuroscience*, vol. 6, no. 6, pp. 449–462, 2005.
- [19] C. Bastian and Y. V. Li, "Fluorescence imaging study of extracellular zinc at the hippocampal mossy fiber synapse," *Neuroscience Letters*, vol. 419, no. 2, pp. 119–124, 2007.
- [20] M. E. Quinta-Ferreira, C. M. Matias, M. Arif, and J. C. Dionísio, "Measurement of presynaptic zinc changes in hippocampal mossy fibers," *Brain Research*, vol. 1026, no. 1, pp. 1–10, 2004.
- [21] P. Paoletti, A. M. Vergnano, B. Barbour, and M. Casado, "Zinc at glutamatergic synapses," *Neuroscience*, vol. 158, no. 1, pp. 126–136, 2009.
- [22] S. Peters, J. Koh, and D. W. Choi, "Zinc selectively blocks the action of N-methyl-D-aspartate on cortical neurons," *Science*, vol. 236, no. 4801, pp. 589–593, 1987.
- [23] G. L. Westbrook and M. L. Mayer, "Micromolar concentrations of Zn^{2+} antagonize NMDA and GABA responses of hippocampal neurons," *Nature*, vol. 328, no. 6131, pp. 640–643, 1987.
- [24] C. W. Christine and D. W. Choi, "Effect of zinc on NMDA receptor-mediated channel currents in cortical neurons," *Journal of Neuroscience*, vol. 10, no. 1, pp. 108–116, 1990.
- [25] P. Legendre and G. L. Westbrook, "The inhibition of single N-methyl-D-aspartate-activated channels by zinc ions on cultured rat neurones," *Journal of Physiology*, vol. 429, pp. 429–449, 1990.
- [26] N. Chen, A. Moshaver, and L. A. Raymond, "Differential sensitivity of recombinant N-methyl-D-aspartate receptor subtypes to zinc inhibition," *Molecular Pharmacology*, vol. 51, no. 6, pp. 1015–1023, 1997.
- [27] P. Paoletti, P. Ascher, and J. Neyton, "High-affinity zinc inhibition of NMDA NR1-NR2A receptors," *Journal of Neuroscience*, vol. 17, no. 15, pp. 5711–5725, 1997.
- [28] J. Y. Koh, "Zinc toxicity on cultured cortical neurons: involvement of N-methyl-D-aspartate receptors," *Neuroscience*, vol. 60, no. 4, pp. 1049–1057, 1994.
- [29] J. H. Weiss, D. M. Hartley, J. Y. Koh, and D. W. Choi, "AMPA receptor activation potentiates zinc neurotoxicity," *Neuron*, vol. 10, no. 1, pp. 43–49, 1993.
- [30] D. D. Lin, A. S. Cohen, and D. A. Coulter, "Zinc-induced augmentation of excitatory synaptic currents and glutamate receptor responses in hippocampal CA3 neurons," *Journal of Neurophysiology*, vol. 85, no. 3, pp. 1185–1196, 2001.
- [31] Y. Jia, J. M. Jeng, S. L. Sensi, and J. H. Weiss, " Zn^{2+} currents are mediated by calcium-permeable AMPA/kainate channels in cultured murine hippocampal neurones," *Journal of Physiology*, vol. 543, no. 1, pp. 35–48, 2002.
- [32] A. Mathie, G. L. Sutton, C. E. Clarke, and E. L. Veale, "Zinc and copper: pharmacological probes and endogenous modulators of neuronal excitability," *Pharmacology and Therapeutics*, vol. 111, no. 3, pp. 567–583, 2006.
- [33] J. Magistretti, L. Castelli, V. Taglietti, and F. Tanzi, "Dual effect of Zn^{2+} on multiple types of voltage-dependent Ca currents in rat palaeocortical neurons," *Neuroscience*, vol. 117, no. 2, pp. 249–264, 2003.
- [34] H. S. Sun, K. Hui, D. W. K. Lee, and Z. P. Feng, " Zn^{2+} sensitivity of high- and low-voltage activated calcium channels," *Biophysical Journal*, vol. 93, no. 4, pp. 1175–1183, 2007.
- [35] K. Inoue, D. Branigan, and Z. G. Xiong, "Zinc-induced neurotoxicity mediated by transient receptor potential melastatin 7 channels," *The Journal of Biological Chemistry*, vol. 285, no. 10, pp. 7430–7439, 2010.
- [36] P. Molnár and J. V. Nadler, "Synaptically-released zinc inhibits N-methyl-D-aspartate receptor activation at recurrent mossy fiber synapses," *Brain Research*, vol. 910, no. 1–2, pp. 205–207, 2001.
- [37] K. Vogt, J. Mellor, G. Tong, and R. Nicoll, "The actions of synaptically released zinc at hippocampal mossy fiber synapses," *Neuron*, vol. 26, no. 1, pp. 187–196, 2000.
- [38] S. Ueno, M. Tsukamoto, T. Hirano et al., "Mossy fiber Zn^{2+} spillover modulates heterosynaptic N-methyl-D-aspartate receptor activity in hippocampal CA3 circuits," *Journal of Cell Biology*, vol. 158, no. 2, pp. 215–220, 2002.
- [39] Y. M. Lu, F. A. Taverna, R. Tu et al., "Endogenous Zn^{2+} is required for the induction of long-term potentiation at rat hippocampal mossy fiber-CA3 synapses," *Synapse*, vol. 38, no. 2, pp. 187–197, 2000.
- [40] A. R. Kay and K. Tóth, "Influence of location of a fluorescent zinc probe in brain slices on its response to synaptic activation," *Journal of Neurophysiology*, vol. 95, no. 3, pp. 1949–1956, 2006.
- [41] A. R. Kay, "Evidence for chelatable zinc in the extracellular space of the hippocampus, but little evidence for synaptic release of Zn ," *Journal of Neuroscience*, vol. 23, no. 17, pp. 6847–6855, 2003.
- [42] S. Yamasaki, K. Sakata-Sogawa, A. Hasegawa et al., "Zinc is a novel intracellular second messenger," *Journal of Cell Biology*, vol. 177, no. 4, pp. 637–645, 2007.
- [43] M. A. Leissring, "The A β Cs of A β -cleaving proteases," *The Journal of Biological Chemistry*, vol. 283, no. 44, pp. 29645–29649, 2008.
- [44] A. I. Bush, G. Multhaup, R. D. Moir et al., "A novel zinc(II) binding site modulates the function of the β A4 amyloid protein precursor of Alzheimer's disease," *The Journal of Biological Chemistry*, vol. 268, no. 22, pp. 16109–16112, 1993.
- [45] A. I. Bush, W. H. Pettingell, M. De Paradis, R. E. Tanzi, and W. Wasco, "The amyloid β -protein precursor and its mammalian homologues. Evidence for a zinc-modulated heparin-binding superfamily," *The Journal of Biological Chemistry*, vol. 269, no. 43, pp. 26618–26621, 1994.
- [46] G. Multhaup, A. I. Busha, P. Pollweina, and C. L. Masters, "Interaction between the zinc(II) and the heparin binding site of the Alzheimer's disease β A4 amyloid precursor protein (APP)," *FEBS Letters*, vol. 355, no. 2, pp. 151–154, 1994.
- [47] G. Multhaup, T. Ruppert, A. Schlicksupp et al., "Copper-binding amyloid precursor protein undergoes a site-specific fragmentation in the reduction of hydrogen peroxide," *Biochemistry*, vol. 37, no. 20, pp. 7224–7230, 1998.
- [48] V. W. Chow, M. P. Mattson, P. C. Wong, and M. Gleichmann, "An overview of APP processing enzymes and products," *Neuromolecular Medicine*, vol. 12, no. 1, pp. 1–12, 2010.
- [49] D. Edbauer, E. Winkler, J. T. Regula, B. Pesold, H. Steiner, and C. Haass, "Reconstitution of γ -secretase activity," *Nature Cell Biology*, vol. 5, no. 5, pp. 486–488, 2003.
- [50] A. Nikolaev, T. McLaughlin, D. D. M. O'Leary, and M. Tessier-Lavigne, "APP binds DR6 to trigger axon pruning and neuron death via distinct caspases," *Nature*, vol. 457, no. 7232, pp. 981–989, 2009.
- [51] Ł. P. Slomnicki and W. Leśniak, "A putative role of the Amyloid Precursor Protein Intracellular Domain (AICD) in transcription," *Acta Neurobiologiae Experimentalis*, vol. 68, no. 2, pp. 219–228, 2008.

- [52] D. R. Edwards, M. M. Handsley, and C. J. Pennington, "The ADAM metalloproteinases," *Molecular Aspects of Medicine*, vol. 29, no. 5, pp. 258–289, 2009.
- [53] W. Bode, F. X. Gomis-Ruth, and W. Stockler, "Astacins, seralysins, snake venom and matrix metalloproteinases exhibit identical zinc-binding environments (HEXXHXXGXXH and Met-turn) and topologies and should be grouped into a common family, the 'metzincins'," *FEBS Letters*, vol. 331, no. 1–2, pp. 134–140, 1993.
- [54] F. X. Gomis-Rüth, "Structural aspects of the metzincin clan of metalloendopeptidases," *Applied Biochemistry and Biotechnology Part B*, vol. 24, no. 2, pp. 157–202, 2003.
- [55] T. M. J. Allinson, E. T. Parkin, A. J. Turner, and N. M. Hooper, "ADAMs family members as amyloid precursor protein alpha-secretases," *Journal of Neuroscience Research*, vol. 74, no. 3, pp. 342–352, 2003.
- [56] M. Asai, C. Hattori, B. Szabó et al., "Putative function of ADAM9, ADAM10, and ADAM17 as APP α -secretase," *Biochemical and Biophysical Research Communications*, vol. 301, no. 1, pp. 231–235, 2003.
- [57] F. Fahrenholz, S. Gilbert, E. Kojro, S. Lammich, and R. Postina, " α -Secretase activity of the disintegrin metalloprotease ADAM 10: influences of domain structure," *Annals of the New York Academy of Sciences*, vol. 920, pp. 215–222, 2000.
- [58] C. Tanabe, N. Hotoda, N. Sasagawa, A. Sehara-Fujisawa, K. Maruyama, and S. Ishiura, "ADAM19 is tightly associated with constitutive Alzheimer's disease APP α -secretase in A172 cells," *Biochemical and Biophysical Research Communications*, vol. 352, no. 1, pp. 111–117, 2007.
- [59] R. Postina, A. Schroeder, I. Dewachter et al., "A disintegrin-metalloproteinase prevents amyloid plaque formation and hippocampal defects in an Alzheimer disease mouse model," *Journal of Clinical Investigation*, vol. 113, no. 10, pp. 1456–1464, 2004.
- [60] D. Hartmann, B. de Strooper, L. Serneels et al., "The disintegrin/metalloprotease ADAM 10 is essential for Notch signalling but not for α -secretase activity in fibroblasts," *Human Molecular Genetics*, vol. 11, no. 21, pp. 2615–2624, 2002.
- [61] P.-H. Kuhn, H. Wang, B. Dislich et al., "ADAM10 is the physiologically relevant, constitutive α -secretase of the amyloid precursor protein in primary neurons," *EMBO Journal*, vol. 29, no. 17, pp. 3020–3032, 2010.
- [62] R. J. Bateman, L. Y. Munsell, J. C. Morris, R. Swarm, K. E. Yarasheski, and D. M. Holtzman, "Human amyloid- β synthesis and clearance rates as measured in cerebrospinal fluid in vivo," *Nature Medicine*, vol. 12, no. 7, pp. 856–861, 2006.
- [63] S. S. El-Amouri, H. Zhu, J. Yu, R. Marr, I. M. Verma, and M. S. Kindy, "Nepriylisin: an enzyme candidate to slow the progression of Alzheimer's disease," *American Journal of Pathology*, vol. 172, no. 5, pp. 1342–1354, 2008.
- [64] L. B. Hersh and D. W. Rodgers, "Nepriylisin and amyloid beta peptide degradation," *Current Alzheimer Research*, vol. 5, no. 2, pp. 225–231, 2008.
- [65] J. S. Miners, S. Baig, J. Palmer, L. E. Palmer, P. G. Kehoe, and S. Love, " $A\beta$ -degrading enzymes in Alzheimer's disease," *Brain Pathology*, vol. 18, no. 2, pp. 240–252, 2008.
- [66] E. R. L. C. Vardy, A. J. Catto, and N. M. Hooper, "Proteolytic mechanisms in amyloid- β metabolism: therapeutic implications for Alzheimer's disease," *Trends in Molecular Medicine*, vol. 11, no. 10, pp. 464–472, 2005.
- [67] H. Kanemitsu, T. Tomiyama, and H. Mori, "Human neprilysin is capable of degrading amyloid β peptide not only in the monomeric form but also the pathological oligomeric form," *Neuroscience Letters*, vol. 350, no. 2, pp. 113–116, 2003.
- [68] J. Apelt, K. Ach, and R. Schliebs, "Aging-related down-regulation of neprilysin, a putative β -amyloid-degrading enzyme, in transgenic Tg2576 Alzheimer-like mouse brain is accompanied by an astroglial upregulation in the vicinity of β -amyloid plaques," *Neuroscience Letters*, vol. 339, no. 3, pp. 183–186, 2003.
- [69] W. Farris, S. G. Schütz, J. R. Cirrito et al., "Loss of neprilysin function promotes amyloid plaque formation and causes cerebral amyloid angiopathy," *American Journal of Pathology*, vol. 171, no. 1, pp. 241–251, 2007.
- [70] H. Akiyama, H. Kondo, K. Ikeda, M. Kato, and P. L. McGeer, "Immunohistochemical localization of neprilysin in the human cerebral cortex: inverse association with vulnerability to amyloid β -protein ($A\beta$) deposition," *Brain Research*, vol. 902, no. 2, pp. 277–281, 2001.
- [71] E. A. Eckman, S. K. Adams, F. J. Troendle et al., "Regulation of steady-state β -amyloid levels in the brain by neprilysin and endothelin-converting enzyme but not angiotensin-converting enzyme," *The Journal of Biological Chemistry*, vol. 281, no. 41, pp. 30471–30478, 2006.
- [72] E. Hellström-Lindahl, R. Ravid, and A. Nordberg, "Age-dependent decline of neprilysin in Alzheimer's disease and normal brain: inverse correlation with $A\beta$ levels," *Neurobiology of Aging*, vol. 29, no. 2, pp. 210–221, 2008.
- [73] W. C. Duckworth, R. G. Bennett, and F. G. Hamel, "A direct inhibitory effect of insulin on a cytosolic proteolytic complex containing insulin-degrading enzyme and multicatalytic proteinase," *The Journal of Biological Chemistry*, vol. 269, no. 40, pp. 24575–24580, 1994.
- [74] K. Vekrellis, Z. Ye, W. Q. Qiu et al., "Neurons regulate extracellular levels of amyloid β -protein via proteolysis by insulin-degrading enzyme," *Journal of Neuroscience*, vol. 20, no. 5, pp. 1657–1665, 2000.
- [75] S. Wang, R. Wang, L. Chen, D. A. Bennett, D. W. Dickson, and D.-S. Wang, "Expression and functional profiling of neprilysin, insulin-degrading enzyme, and endothelin-converting enzyme in prospectively studied elderly and Alzheimer's brain," *Journal of Neurochemistry*, vol. 115, no. 1, pp. 47–57, 2010.
- [76] M. C. Leal, V. B. Dorfman, A. F. Gamba et al., "Plaque-associated overexpression of insulin-degrading enzyme in the cerebral cortex of aged transgenic Tg2576 mice with Alzheimer pathology," *Journal of Neuropathology and Experimental Neurology*, vol. 65, no. 10, pp. 976–987, 2006.
- [77] B. C. Miller, E. A. Eckman, K. Sambamurti et al., "Amyloid- β peptide levels in brain are inversely correlated with insulin activity levels in vivo," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 10, pp. 6221–6226, 2003.
- [78] A. P. Davenport, R. E. Kuc, and J. W. Mockridge, "Endothelin-converting enzyme in the human vasculature: evidence for differential conversion of big endothelin-3 by endothelial and smooth-muscle cells," *Journal of Cardiovascular Pharmacology*, vol. 31, supplement 1, pp. S1–S3, 1998.
- [79] K. J. Yin, J. R. Cirrito, P. Yan et al., "Matrix metalloproteinases expressed by astrocytes mediate extracellular amyloid- β peptide catabolism," *Journal of Neuroscience*, vol. 26, no. 43, pp. 10939–10948, 2006.
- [80] A. Falkevall, N. Alikhani, S. Bhushan et al., "Degradation of the amyloid β -protein by the novel mitochondrial

- peptidasome, PreP," *The Journal of Biological Chemistry*, vol. 281, no. 39, pp. 29096–29104, 2006.
- [81] M. A. Lovell, J. D. Robertson, W. J. Teesdale, J. L. Campbell, and W. R. Markesbery, "Copper, iron and zinc in Alzheimer's disease senile plaques," *Journal of the Neurological Sciences*, vol. 158, no. 1, pp. 47–52, 1998.
- [82] J. Y. Lee, T. B. Cole, R. D. Palmiter, S. W. Suh, and J. Y. Koh, "Contribution by synaptic zinc to the gender-disparate plaque formation in human Swedish mutant APP transgenic mice," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 11, pp. 7705–7710, 2002.
- [83] A. I. Bush, W. H. Pettingell, G. Multhaup et al., "Rapid induction of Alzheimer A β amyloid formation by zinc," *Science*, vol. 265, no. 5177, pp. 1464–1467, 1994.
- [84] K. H. Lim, Y. K. Kim, and Y. T. Chang, "Investigations of the molecular mechanism of metal-induced A β (1–40) amyloidogenesis," *Biochemistry*, vol. 46, no. 47, pp. 13523–13532, 2007.
- [85] L. M. Miller, Q. Wang, T. P. Telivala, R. J. Smith, A. Lanzirotti, and J. Miklossy, "Synchrotron-based infrared and X-ray imaging shows focalized accumulation of Cu and Zn co-localized with β -amyloid deposits in Alzheimer's disease," *Journal of Structural Biology*, vol. 155, no. 1, pp. 30–37, 2006.
- [86] S. A. Kozin, S. Zirah, S. Rebuffat, G. Hui Bon Hoa, and P. Debey, "Zinc binding to Alzheimer's A β (1-16) peptide results in stable soluble complex," *Biochemical and Biophysical Research Communications*, vol. 285, no. 4, pp. 959–964, 2001.
- [87] Y. Mekmouche, Y. Coppel, K. Hochgräfe et al., "Characterization of the Zn^{II} binding to the peptide amyloid- β ¹⁻¹⁶ linked to Alzheimer's disease," *ChemBioChem*, vol. 6, no. 9, pp. 1663–1671, 2005.
- [88] C. D. Syme and J. H. Viles, "Solution NMR investigation of Zn²⁺ and Cd²⁺ binding to amyloid-beta peptide (A β) of Alzheimer's disease," *Biochimica et Biophysica Acta*, vol. 1764, no. 2, pp. 246–256, 2006.
- [89] C. Talmard, A. Bouzan, and P. Faller, "Zinc binding to amyloid- β : isothermal titration calorimetry and Zn competition experiments with Zn sensors," *Biochemistry*, vol. 46, no. 47, pp. 13658–13666, 2007.
- [90] S. Zirah, S. A. Kozin, A. K. Mazur et al., "Structural changes of region 1-16 of the Alzheimer disease amyloid β -peptide upon zinc binding and in vitro aging," *The Journal of Biological Chemistry*, vol. 281, no. 4, pp. 2151–2161, 2006.
- [91] P. Faller and C. Hureau, "Bioinorganic chemistry of copper and zinc ions coordinated to amyloid- β peptide," *Dalton Transactions*, no. 7, pp. 1080–1094, 2009.
- [92] C. Talmard, L. Guilloreau, Y. Coppel, H. Mazarguil, and P. Faller, "Amyloid-beta peptide forms monomeric complexes with Cu^{II} and Zn^{II} prior to aggregation," *ChemBioChem*, vol. 8, no. 2, pp. 163–165, 2007.
- [93] J. Danielsson, R. Pierattelli, L. Banci, and A. Gräslund, "High-resolution NMR studies of the zinc-binding site of the Alzheimer's amyloid β -peptide," *FEBS Journal*, vol. 274, no. 1, pp. 46–59, 2007.
- [94] S. Zirah, S. Rebuffat, S. A. Kozin et al., "Zinc binding properties of the amyloid fragment A β (1–16) studied by electrospray-ionization mass spectrometry," *International Journal of Mass Spectrometry*, vol. 228, no. 2-3, pp. 999–1016, 2003.
- [95] C. C. Curtain, F. Ali, I. Volitakis et al., "Alzheimer's disease amyloid-beta binds copper and zinc to generate an allosterically ordered membrane-penetrating structure containing superoxide dismutase-like subunits," *The Journal of Biological Chemistry*, vol. 276, no. 23, pp. 20466–20473, 2001.
- [96] S. T. Liu, G. Howlett, and C. J. Barrow, "Histidine-13 is a crucial residue in the zinc ion-induced aggregation of the A β peptide of Alzheimer's disease," *Biochemistry*, vol. 38, no. 29, pp. 9373–9378, 1999.
- [97] D. S. Yang, J. McLaurin, K. Qin, D. Westaway, and P. E. Fraser, "Examining the zinc binding site of the amyloid- β peptide," *European Journal of Biochemistry*, vol. 267, no. 22, pp. 6692–6698, 2000.
- [98] J. Dong, C. S. Atwood, V. E. Anderson et al., "Metal binding and oxidation of amyloid- β within isolated senile plaque cores: raman microscopic evidence," *Biochemistry*, vol. 42, no. 10, pp. 2768–2773, 2003.
- [99] W. Garzon-Rodriguez, A. K. Yatsimirsky, and C. G. Glabe, "Binding of Zn(II), Cu(II), and Fe(II) ions to Alzheimer's A β peptide studied by fluorescence," *Bioorganic and Medicinal Chemistry Letters*, vol. 9, no. 15, pp. 2243–2248, 1999.
- [100] F. Ricchelli, D. Drago, B. Filippi, G. Tognon, and P. Zatta, "Aluminum-triggered structural modifications and aggregation of β -amyloids," *Cellular and Molecular Life Sciences*, vol. 62, no. 15, pp. 1724–1733, 2005.
- [101] V. Töugu, A. Karafin, and P. Palumaa, "Binding of zinc(II) and copper(II) to the full-length Alzheimer's amyloid- β peptide," *Journal of Neurochemistry*, vol. 104, no. 5, pp. 1249–1259, 2008.
- [102] L. L. Rulon, J. D. Robertson, M. A. Lovell, M. A. Deibel, W. D. Ehmann, and W. R. Markesbery, "Serum zinc levels and Alzheimer's disease," *Biological Trace Element Research*, vol. 75, no. 1–3, pp. 79–85, 2000.
- [103] F. J. Jiménez-Jiménez, J. A. Molina, M. V. Aguilar et al., "Cerebrospinal fluid levels of transition metals in patients with Parkinson's disease," *Journal of Neural Transmission*, vol. 105, no. 4-5, pp. 497–505, 1998.
- [104] C. Jeandel, M. B. Nicolas, F. Dubois, F. Nabet-Belleville, F. Penin, and G. Cuny, "Lipid peroxidation and free radical scavengers in Alzheimer's disease," *Gerontology*, vol. 35, no. 5-6, pp. 275–282, 1989.
- [105] L. Baum, I. H. S. Chan, S. K. K. Cheung et al., "Serum zinc is decreased in Alzheimer's disease and serum arsenic correlates positively with cognitive ability," *BioMetals*, vol. 23, no. 1, pp. 173–179, 2010.
- [106] M. A. Lovell, J. L. Smith, S. Xiong, and W. R. Markesbery, "Alterations in zinc transporter protein-1 (ZnT-1) in the brain of subjects with mild cognitive impairment, early, and late-stage Alzheimer's disease," *Neurotoxicity Research*, vol. 7, no. 4, pp. 265–271, 2005.
- [107] J. L. Smith, S. Xiong, W. R. Markesbery, and M. A. Lovell, "Altered expression of zinc transporters-4 and -6 in mild cognitive impairment, early and late Alzheimer's disease brain," *Neuroscience*, vol. 140, no. 3, pp. 879–888, 2006.
- [108] M. A. Lovell, "A potential role for alterations of zinc and zinc transport proteins in the progression of Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 16, no. 3, pp. 471–483, 2009.
- [109] M. A. Smith, P. L. R. Harris, L. M. Sayre, and G. Perry, "Iron accumulation in Alzheimer disease is a source of redox-generated free radicals," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 18, pp. 9866–9868, 1997.
- [110] D. A. Butterfield, T. Reed, S. F. Newman, and R. Sultana, "Roles of amyloid β -peptide-associated oxidative stress and brain protein modifications in the pathogenesis of Alzheimer's disease and mild cognitive impairment," *Free Radical Biology and Medicine*, vol. 43, no. 5, pp. 658–677, 2007.

- [111] A. Clements, D. Allsop, D. M. Walsh, and C. H. Williams, "Aggregation and metal-binding properties of mutant forms of the amyloid A β peptide of Alzheimer's disease," *Journal of Neurochemistry*, vol. 66, no. 2, pp. 740–747, 1996.
- [112] W. P. Esler, E. R. Stimson, J. M. Jennings, J. R. Ghilardi, P. W. Mantyh, and J. E. Maggio, "Zinc-induced aggregation of human and rat β -amyloid peptides in vitro," *Journal of Neurochemistry*, vol. 66, no. 2, pp. 723–732, 1996.
- [113] J. Dong, J. E. Shokes, R. A. Scott, and D. G. Lynn, "Modulating amyloid self-assembly and fibril morphology with Zn(II)," *Journal of the American Chemical Society*, vol. 128, no. 11, pp. 3540–3542, 2006.
- [114] Y. Miller, B. Ma, and R. Nussinov, "Zinc ions promote Alzheimer $\alpha\beta$ aggregation via population shift of polymorphic states," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 21, pp. 9490–9495, 2010.
- [115] A. Deshpande, H. Kawai, R. Metherate, C. G. Glabe, and J. Busciglio, "A role for synaptic zinc in activity-dependent $\alpha\beta$ oligomer formation and accumulation at excitatory synapses," *Journal of Neuroscience*, vol. 29, no. 13, pp. 4004–4015, 2009.
- [116] R. A. Cherny, J. T. Legg, C. A. McLean et al., "Aqueous dissolution of Alzheimer's disease A β amyloid deposits by biometal depletion," *The Journal of Biological Chemistry*, vol. 274, no. 33, pp. 23223–23228, 1999.
- [117] R. A. Cherny, C. S. Atwood, M. E. Xilinas et al., "Treatment with a copper-zinc chelator markedly and rapidly inhibits β -amyloid accumulation in Alzheimer's disease transgenic mice," *Neuron*, vol. 30, no. 3, pp. 665–676, 2001.
- [118] C. Opazo, S. Luza, V. L. Villemagne et al., "Radioiodinated clioquinol as a biomarker for β -amyloid: Zn²⁺ complexes in Alzheimer's disease," *Aging Cell*, vol. 5, no. 1, pp. 69–79, 2006.
- [119] C. W. Ritchie, A. I. Bush, A. Mackinnon et al., "Metal-protein attenuation with iodochlorhydroxyquin (clioquinol) targeting Abeta amyloid deposition and toxicity in Alzheimer disease: a pilot phase 2 clinical trial," *Archives of Neurology*, vol. 60, no. 12, pp. 1685–1691, 2003.
- [120] A. M. Mancino, S. S. Hindo, A. Kochi, and M. H. Lim, "Effects of clioquinol on metal-triggered amyloid-beta aggregation revisited," *Inorganic Chemistry*, vol. 48, no. 20, pp. 9596–9598, 2009.
- [121] L. Lannfelt, K. Blennow, H. Zetterberg et al., "Safety, efficacy, and biomarker findings of PBT2 in targeting A β as a modifying therapy for Alzheimer's disease: a phase IIa, double-blind, randomised, placebo-controlled trial," *The Lancet Neurology*, vol. 7, no. 9, pp. 779–786, 2008.
- [122] Z. Y. Mo, Y. Z. Zhu, H. L. Zhu, J. B. Fan, J. Chen, and Y. Liang, "Low micromolar zinc accelerates the fibrillization of human Tau via bridging of Cys-291 and Cys-322," *The Journal of Biological Chemistry*, vol. 284, no. 50, pp. 34648–34657, 2009.
- [123] W. L. An, C. Bjorkdahl, R. Liu, R. F. Cowburn, B. Winblad, and J. J. Pei, "Mechanism of zinc-induced phosphorylation of p70 S6 kinase and glycogen synthase kinase 3 β in SH-SY5Y neuroblastoma cells," *Journal of Neurochemistry*, vol. 92, no. 5, pp. 1104–1115, 2005.
- [124] T. B. Cole, A. Martyanova, and R. D. Palmiter, "Removing zinc from synaptic vesicles does not impair spatial learning, memory, or sensorimotor functions in the mouse," *Brain Research*, vol. 891, no. 1-2, pp. 253–265, 2001.
- [125] P. A. Adlard, J. M. Parncutt, D. I. Finkelstein, and A. I. Bush, "Cognitive loss in zinc transporter-3 knock-out mice: a phenocopy for the synaptic and memory deficits of Alzheimer's disease?" *Journal of Neuroscience*, vol. 30, no. 5, pp. 1631–1636, 2010.
- [126] G. Schmitt-Ulms, S. Ehsani, J. C. Watts, D. Westaway, and H. Wille, "Evolutionary descent of prion genes from the ZIP family of metal Ion transporters," *PLoS ONE*, vol. 4, no. 9, Article ID e7208, 2009.
- [127] J. A. Duce, A. Tsatsanis, M. A. Cater et al., "Iron-export ferroxidase activity of beta-amyloid precursor protein is inhibited by zinc in Alzheimer's disease," *Cell*, vol. 142, no. 6, pp. 857–867, 2010.
- [128] I. J. Whitehouse, C. Jackson, A. J. Turner, and N. M. Hooper, "Prion protein is reduced in aging and in sporadic but not in familial Alzheimer's Disease," *The Journal of Alzheimer's Disease*, vol. 22, pp. 1023–1031, 2010.
- [129] E. T. Parkin, N. T. Watt, I. Hussain et al., "Cellular prion protein regulates β -secretase cleavage of the Alzheimer's amyloid precursor protein," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 26, pp. 11062–11067, 2007.
- [130] N. T. Watt, D. R. Taylor, A. Gillott, D. A. Thomas, W. S. S. Perera, and N. M. Hooper, "Reactive oxygen species-mediated β -cleavage of the prion protein in the cellular response to oxidative stress," *The Journal of Biological Chemistry*, vol. 280, no. 43, pp. 35914–35921, 2005.
- [131] C. L. Haigh, S. C. Drew, M. P. Boland et al., "Dominant roles of the polybasic proline motif and copper in the PrP23-89-mediated stress protection response," *Journal of Cell Science*, vol. 122, no. 10, pp. 1518–1528, 2009.